

Letters to the Editor

The Board of Editors does not hold itself responsible for opinions expressed in the letters published in this section. The notes containing short reports of original investigations communicated to this section should not contain many figures and should not exceed 500 words in length. The contributions reaching the Secretary by the 15th of any month may be expected to appear in the issue for the next month. No proof will be sent to the author.

1

MOLECULAR ORBITAL THEORY OF TRIGONALLY DISTORTED $[\text{Co}, 6\text{H}_2\text{O}^{2+}]$ COMPLEX

L. C. JACKSON* AND R. RAI

DEPARTMENT OF MAGNETISM, INDIAN ASSOCIATION FOR THE CULTIVATION OF SCIENCE,
CALCUTTA-32

(Received January 9, 1964)

Under an octahedral field of type O_h , $3d^7$ 4F ground state of Co^{2+} ion splits up into two triplets $^4T_{1g}$, $^4T_{2g}$ and a singlet $^4A_{2g}$ of successively increasing energies. There is another term $^1P(^4T_{1g})$ arising from the same configuration $3d^7$ of the free ion, which lies about $20,000 \text{ cm}^{-1}$ above $^4T_{1g}$ in crystals (Abragam *et al*, 1951). So that the lowest triplet state contains an admixture of 4P through the effective orbital Lande g -factors α , α' (Bose *et al*, 1961), which are appreciably different in crystals from the value $3/2$ for the free ion F -state. For trigonal distortion of the octahedron, the appropriate Hamiltonian is given by

$$H = V_{\text{trig}} + \alpha u_{\xi} s_{\xi} + \alpha' (u_{\xi} s_{\xi} + u_{\eta} s_{\eta})$$

where the effect of Spin-orbit interaction takes the form $\Sigma(us)_i$ (Bose *et al*, 1960). Operating above Hamiltonian over the appropriate trigonal orbital states for the




Fig. 1

* Royal Military College, Canada, Kingston, Ontario, Canada.

lowest triplet (Bose *et al*, 1963) formed by mixing of the central *d*- and surrounding *p*-orbitals, we get the energies and wave-functions for lowest six Kramer's doublets, ready for magnetic perturbation.

Thus we have derived a very complicated expression for the principal ionic magnetic susceptibilities K_i ($i = \parallel$ or \perp) along and perpendicular to the trigonal axis of the $\text{CoSiF}_6 \cdot 6\text{H}_2\text{O}$ single crystal, which is identical for the single trigonally distorted $[\text{Co}^{2+} \cdot 6\text{H}_2\text{O}]$ complex in the unit cell sp.gr C_{3i} (Pauling 1930) on the basis of molecular orbital theory of Stevens (1953), Bose *et al*, (1960) and compared our theoretical with the experimental results by one of us (L.C.J.).

We had to increase the trigonal field coefficient Δ with temperature, from a value of 830 cm^{-1} at 1.67°K to 952 cm^{-1} at 90°K and then decrease it to 560 cm^{-1} at 290.4°K ; the values for the spin-orbit coupling coefficient we have to take $\zeta_{\parallel} = -137 \text{ cm}^{-1}$ and $\zeta_{\perp} = -130 \text{ cm}^{-1}$ instead of free ion value -180 cm^{-1} ; the other parameters—the effective orbital Lande factors $\alpha = 1.242$; $\alpha' = 1.495$ and orbital reduction factors $k_{\parallel} = .975$ $k_{\perp} = .93$.

The anisotropic reduction in spin-orbit coupling coefficient is due to overlap of the Co^{2+} charge clouds with *s* and *p* ligand charge clouds. The increase in Δ below 90°K is, as observed earlier, due to thermal expansion or relaxation effects but the decrease above 90°K appears to be due to some kind of phase transition reversible in character, evidence for which is available from some of our recent measurements between 90°K and 300°K (Mazumdar *et al*). Details of the theoretical developments and experimental results will be published elsewhere.

ACKNOWLEDGMENT

The junior author (R. R.) is indebted to Prof. A. Bose, D.Sc., F.N.I. for suggesting the problem and guidance in the work. Many valuable discussions and suggestions by Dr. R. Chatterjee are also gratefully acknowledged.

REFERENCES

- Abragram, A. and Pryce, M. H. L., 1951, *Proc. Roy. Soc. A.*, **206**, 173.
- Bose, A., Chakravarty, A. S. and Chatterjee R., 1961, *Proc. Roy. Soc., A*, **261**, 43.
- Bose, A., Chakravarty, A. S. and Chatterjee, R., 1960, *Proc. Roy. Soc.* **255**, 145.
- Bose, A., Chatterjee, R. and Rai, R., 1963, *Proc. phys. Soc.* (communicated),
- Mazumdar, M. and Dutta, S. K., *Jour. Chem. Phys.* (Communicated)
- Pauling, L. C., 1930, *Zeits. F. Kristallograhie* **72A**, 482.
- Stevens, K. W. H., 1953, *Proc. Roy. Soc., A* **219**, 542.